

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 September 2005 (22.09.2005)

PCT

(10) International Publication Number
WO 2005/087950 A1

(51) International Patent Classification⁷: **C12Q 1/68**

(21) International Application Number:
PCT/JP2005/004881

(22) International Filing Date: 14 March 2005 (14.03.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2004-070986 12 March 2004 (12.03.2004) JP

(71) Applicant (for all designated States except US): **CANON KABUSHIKI KAISHA** [JP/JP]; 3-30-2, Shimomaruko, Ohta-ku, Tokyo, 1468501 (JP).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **OKAMOTO, Tadashi** [JP/JP]; c/o CANON KABUSHIKI KAISHA, 3-30-2, Shimomaruko, Ohta-ku, Tokyo, 1468501 (JP).

(74) Agents: **OKABE, Masao** et al.; No. 602, Fuji Bldg., 2-3, Marunouchi 3-chome, Chiyoda-ku, Tokyo 1000005 (JP).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF DETECTING NUCLEIC ACID USING AMPLIFICATION ON AN ARRAY

(57) Abstract: Provided is a method of detecting a nucleic acid, which enables simple, high-efficiency, and high-precise detection of various genes and may be widely utilized in the fields on the basis of gene detection. Provided is a method of detecting a nucleic acid, which enables a solid-phase universal PCR, solid-phase multiplex PCR, or solid-phase universal multiplex PCR using a nucleic acid array.



WO 2005/087950 A1

DESCRIPTION

METHOD OF DETECTING NUCLEIC ACID USING AMPLIFICATION ON AN ARRAY

5 TECHNICAL FIELD

The present invention relates to a method of detecting a nucleic acid, and more particularly to a method of detecting a nucleic acid using a primer array and a PCR method.

10

BACKGROUND ART

Base sequence analysis of genome or the like of various living beings including human beings has intensively proceeded, with the result that gene analysis has come into use for diagnosis of a genetic disease, cancer, infectious disease, lifestyle-related disease, or the like; decision of a therapeutic strategy; check after treatment; and prognostic expectation. Moreover, in another field, gene analysis is beginning to be used for distribution management of foods such as meat and grain.

20

Among techniques used for the gene analysis, a nucleic acid array immobilized with plural nucleic acids such as DNA probes, oligonucleotide probes, and cDNA is one of the techniques that has particularly drawn attention since the late 1990s. The nucleic

25

acid array may also be referred to as DNA array,
oligonucleotide array, cDNA array, or the like, and
in some situations, as nucleic acid chip or DNA chip.
Gene analysis techniques using a nucleic acid array
5 or a nucleic acid array itself essentially have an
advantage that multiple and various solid-phase
hybridizations can be performed simultaneously,
although which is not described in detail herein
because those techniques have been generally well
10 known recently.

On the other hand, the PCR (polymerase chain
reaction) method that has been considered as an
important technique for gene analysis does not go out
of fashion today and is one technique most widely
15 used in the gene analysis.

Since the PCR method was devised, some
improvements and evolutions have been accomplished.
For example, the traditional PCR method was performed
for one gene or one base sequence in the form that a
20 base sequence to be amplified was sandwiched by two
primers. However, recently, PCR is performed using
two primers sandwiching base sequences to be
amplified that include different base sequences of
plural genes or plural base sequences. That is,
25 amplification of plural genes or the like can be
realized using only two (common) primers. Such PCR
method is generally referred to as a universal PCR

method. For example, when plural infecting organisms causing infectious diseases are to be identified, PCR is performed using primers which sandwich base sequences including a part of base sequences of a specific rRNA (for example, 16s rRNA) unique to the respective infecting organisms and have common base sequences in the respective infecting organisms, and the aforementioned unique base sequences are detected, to thereby identify and detect the respective infecting organisms. Such method has an advantage that PCR procedures are not required for the respective plural genes or the like.

On the other hand, a procedure called multiplex PCR has been developed. This procedure is intended to amplify plural genes or the like with different primer sets in one batch. Although this procedure has a difficulty in design, setting of a primer strand length, base sequence, and PCR conditions or the like, basically, plural and multiple PCR procedures can be performed simultaneously. Moreover, in some situations, improved quantitative detection can be expected through the reactions in the same batch.

Further, a procedure that is a combination of the above-described chip technique and PCR method has also been proposed.

There has been disclosed one procedure of so-

called solid phase PCR that is used for performing plural PCR procedures on one solid-phase by immobilizing a set of PCR primers including two oligonucleotides on a matrix in a nucleic acid array (see Japanese Patent Application Laid-Open No. 2001-299346). According to such a procedure, two amplified products that have been amplified on the matrix form an inverted U-shaped hybridized product mutually by keeping the products under hybridization conditions together with a nucleic acid array after PCR. A gene to be detected can be identified and detected by detecting the hybridized product using, for example, a fluorescent intercalator dye.

Moreover, a solid-phase PCR method that is performed on a microplate having plural wells has been proposed (see THCH NOTE Vol. 3, No. 16, published by Nalge Nunc International). That is, the method of PCR in which, one primer of a set of PCR primers is covalently immobilized on the well surface of the microplate, and a template nucleic acid and both primers are used in the wells for PCR.

Detection is performed separately using a detection probe for PCR amplified products that are elongated from the primer immobilized on the well surfaces.

Conventional techniques relating to the present invention, that is, gene detection techniques, nucleic acid array techniques, PCR methods, and

solid-phase PCR methods have been outlined above.

The aforementioned solid-phase PCR is basically used for one PCR reaction on one matrix or one well. and it is difficult to be applied to the above-

5 described universal PCR or multiplex PCR. Moreover, in the method described in Japanese Patent

Application Laid-Open No. 2001-299346, hybridization is sterically limited because an inverted U-shaped hybridized product is finally formed in a microregion.

10 Moreover, because the reactions of the aforementioned method described in TECH NOTE Vol. 3, No. 16 are performed in microplate wells, the numbers of the reactions are limited compared to reactions in the case of a nucleic acid array, and a large volume of a
15 reaction solution is required, so that the detection efficiency may need more improvement. Further, the detection in those techniques requires another step after amplification, which is performed using, for example, a fluorescent intercalator or another
20 detection probe.

Moreover, Japanese Patent Application Laid-Open No. 2003-523183 discloses an array for detecting and comparing expression patterns of plural target polynucleotides from at least two different
25 biological sources, which includes at least two groups of oligonucleotide primers immobilized on a discrete region of a solid-phase support. Each group

of oligonucleotide primers is selected for a specific target polynucleotide and includes a sequence complementary to the sequence of the specific target polynucleotide. In the array, each group can be
5 identified by the position on the solid-phase support. Further, the array can be used for detecting expression patterns of target polynucleotides from single biological source.

However, in order to detect different genes,
10 only one set of primers is prepared for each detection. In the case where different microorganisms each having extremely similar sequence, for example, infecting organisms of infectious diseases are identified, determination may not be
15 performed.

DISCLOSURE OF THE INVENTION

In view of the above-described circumstances, the present invention relates to a solid-phase PCR
20 method which enables simple, high-efficiency, and high-precise detection of various genes, and an object of the present invention is to provide a method of detecting a nucleic acid which may be widely utilized in fields on the basis of gene
25 detection such as diagnosis, treatment, or prognostic expectation in the medical field; or distribution management in the food field.

The aforementioned object is accomplished by the following aspects of the present invention.

More specifically, a first aspect of the present invention relates to a method of detecting a nucleic acid characterized by including the steps of:

(1) preparing a single-stranded nucleic acid having plural partial and sequential base sequences to be detected (A-strand) and a single-stranded nucleic acid having a base sequence complementary to a base sequence of the A-strand (B-strand);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions;

(3) preparing a nucleic acid having a sequence complementary to a partial and sequential base sequence within the region between a 3'-end of the A-strand and the base sequence to be detected which is located nearest the 3'-end as a primer for elongating the B-strand;

(4) performing PCR reactions using the A-strand and B-strand as templates, and using the primers immobilized on the substrate, and the primer for elongating the B-strand;

(5) forming a hybridized product of a nucleic acid corresponding to the A-strand which has been elongated and amplified as a result of the PCR reactions and bound to the substrate and a nucleic acid corresponding to the B-strand which has been elongated and amplified and has not bound to the substrate; and

(6) detecting the base sequence to be detected by detecting the hybridized product in the respective primer-immobilized regions in the array.

The aforementioned method enables the universal PCR on a nucleic acid array.

A second aspect of the present invention relates to a method of detecting a nucleic acid characterized by including the steps of:

(1) preparing plural single-stranded nucleic acids each having a partial and sequential base sequence to be detected (A-strand group: A1-strand to An-strand: $n \geq 2$) and a group of single-stranded nucleic acids each having a base sequence complementary to a base sequence of each strand of the A-strand group (B-strand group: B1-strand to Bn-strand: $n \geq 2$);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and

preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions;

(3) preparing nucleic acids having a sequence
5 complementary to a partial and sequential base sequence within the region between a 3'-end of each strand of the A-strand group and the base sequence to be detected which is located nearest the 3'-end as primers for elongating the B-strands (PB-strand
10 group: PB1-strand to PBn-strand $n \geq 2$);

(4) performing PCR reactions using each strand of the A-strand group and each strand of B-strand group as templates, and using the primers immobilized on the substrate, and the plural primers for
15 elongating the B-strands of the PB-strand group;

(5) forming a hybridized product of a nucleic acid corresponding to the A-strand group which has been elongated and amplified as a result of the PCR reactions and bound to the substrate and a nucleic
20 acid corresponding to the B-strand group which has been elongated and amplified and has not bound to the substrate; and

(6) detecting the base sequence to be detected by detecting the hybridized product in the respective
25 primer-immobilized regions in the array.

In the aforementioned detection method, each A-strand may include one or more of the plural base

sequences to be detected.

The above-described methods enable the solid-phase universal PCR, solid-phase multiplex PCR and solid-phase universal multiplex PCR using the nucleic acid array, which have not been accomplished by conventional techniques.

Further, a third aspect of the present invention relates to a method of detecting a nucleic acid characterized by including the steps of:

(1) preparing a single-stranded nucleic acid having plural partial and sequential base sequences to be detected (A-strand) and a single-stranded nucleic acid having a base sequence complementary to a base sequence of the A-strand (B-strand);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions;

(3) preparing a nucleic acid having a sequence complementary to a partial and sequential base sequence within the region between a 3'-end of the A-strand and the base sequence to be detected which is located nearest the 3'-end as a primer for elongating the B-strand;

(4) performing PCR reactions using the A-strand and the B-strand as templates, and using the primers immobilized on the substrate, and the primer for elongating the B-strand, and nucleotide monomers with
5 a part or all of at least one of the nucleotide monomers being labeled; and

(5) detecting a nucleic acid corresponding to the A-strand which has been elongated and amplified from a probe binding to the substrate via the label
10 incorporated in the nucleic acid.

Further, a forth aspect of the present invention relates to a method of detecting a nucleic acid characterized by including the steps of:

(1) preparing plural single-stranded nucleic
15 acids each having a partial and sequential base sequence to be detected (A-strand group: A1-strand to An-strand: $n \geq 2$) and a group of single-stranded nucleic acids each having a base sequence complementary to a base sequence of each strand of
20 the A-strand group (B-strand group: B1-strand to Bn-strand: $n \geq 2$);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers
25 independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the

primer-immobilized regions;

(3) preparing nucleic acids each having a sequence complementary to a partial and sequential base sequence within a region between a 3'-end of each strand of the A-strand group and the base sequence to be detected which is located nearest the 3'-end as primers for elongating the B-strands (PB-strand group: PB1-strand to PBn-strand: $n \geq 2$);

(4) performing PCR reactions using each strand of the A-strand group and each strand of the B-strand group as templates, and using the primers immobilized on the substrate and the plural primers for elongating the B-strands of the PB-strand group and nucleotide monomers with a part or all of at least one of the nucleotide monomers being labeled; and

(5) detecting a nucleic acid corresponding to the A-strand which has been elongated and amplified from a probe binding to the substrate via the label incorporated in the nucleic acid.

In the aforementioned detection method, each A-strand may include one or more of the plural base sequences to be detected.

One feature of the third and forth aspects of the present invention is to utilize a labeled monomer as a substitute for a part or all of at least one of nucleotide monomers used in PCR reactions in order to simplify a detection step. The method according to

the aspects enables the solid-phase PCR capable of performing detection immediately after PCR, solid-phase universal PCR capable of performing detection immediately after PCR, solid-phase multiplex PCR, or
5 solid-phase universal multiplex PCR without performing two-tiered steps in which hybridization is performed after PCR.

Further, the method according to the first to forth aspects of the present invention is
10 characterized in that a nucleic acid having the same base sequence as one selected from a region nearer the 5'-end of A-strand than a base sequence to be detected which is nearest the 5'-end (A-strand elongating primer) is added to the B-strand
15 elongating primer, and the nucleic acid is used as a primer which is not immobilized on a substrate, so that the amplifying efficiency of a target base sequence in PCR can be improved.

Moreover, in a further aspect of the method
20 according to the first to forth aspects of the present invention, the method is characterized in that PCR reactions and detection are performed in the form in which primer arrays are present in the same container. Further, the PCR reactions and detection
25 can be performed while being observed continuously using the same means.

Another aspect of the present invention relates

to a detection apparatus which enables the detection method of the present invention, characterized by including: at least a PCR reaction container; and detection means.

5 Moreover, the present invention includes provision of a kit for detecting a nucleic acid, including: the primer arrays; a PCR reaction reagent; and a nucleic acid detecting reagent.

10 Note that reference symbol "n" in the methods according to the respective aspects represents the same number (integer).

 The present invention enables the solid-phase universal PCR, solid-phase multiplex PCR, or solid-phase universal multiplex PCR using nucleic acid
15 arrays. Moreover, the present invention enables, by using a labeled nucleotide monomer in PCR, the solid-phase PCR capable of performing detection immediately after PCR, solid-phase universal PCR capable of performing detection immediately after PCR, solid-
20 phase multiplex PCR, or solid-phase universal multiplex PCR.

 That is, according to the present invention, simple, high-efficiency, and high-precise detection of various and multiple genes can be performed.

25 Other features and advantages of the present invention will be apparent from the following description taken in conjunction with the

accompanying drawings, in which like reference characters designate the same or similar parts throughout the figures thereof.

5 BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a schematic view for illustrating the solid-phase universal PCR according to the present invention; and

10 Fig. 2 is a schematic view for illustrating the solid-phase universal PCR and incorporated label according to the present invention.

Fig. 3 schematically illustrates an apparatus for carrying out the method according to the invention.

15 Fig. 4 schematically illustrates another apparatus for carrying out the method according to the invention.

Fig. 5 schematically illustrates still another apparatus for carrying out the method according to
20 the invention.

BEST MODE FOR CARRYING OUT THE INVENTION

Preferred embodiments of the present invention will now be described in detail in accordance with
25 the accompanying drawings.

A method of detecting a nucleic acid according to a first mode of the present invention includes the

aforementioned steps (1) to (6). This method enables the universal PCR on a primer array.

In this case, if the base sequence to be detected is, for example, of a gene having a double-stranded nucleic acid, the sense base sequence is
5 basically the same as the anti-sense base sequence. It is obvious that analysis of the anti-sense sequence reveals the sense base sequence complementary to the sequence (that is, the base
10 sequence of the gene).

Each operation in each step can be performed using the known method. In the detection step as the final step, any procedure which is capable of detecting a formed hybridized product may be used.
15 For example, a detection method using a fluorescent intercalator dye or fluorescent groove binder dye which interacts with a double-stranded nucleic acid to emit or enhance fluorescence may be utilized. Alternatively, there may also be adopted a procedure
20 for labeling a nucleic acid having a base sequence complementary to a base sequence located in the position nearer 3'-end than that of the 3'-end of the base sequence to be detected of A-strand, which is used as a primer in the step (4) (which is a B-strand
25 elongating primer and a primer derived from a segment nearer the 3'-end of A-strand; herein, the phrase "derived from a position nearer the 3'-end" means

"derived from a segment located in the position nearer 3'-end than that of the base sequence to be detected which is nearest the 3'-end"), with a fluorescent dye such as fluorescein, tetramethylrhodamine, Cy3, or Cy5, or radioisotope. Note that, in the case of using the fluorescent dye, a fluorescent microscope may be used for observation, and a step of observing a label using, for example, a confocal fluorescent microscope may further be included.

In this method, one primer of a set of primers to be used PCR reactions is immobilized on a solid-phase, and the other (B-strand elongating primer) is dissolved in a reaction solution before use. The B-strand elongating primer has functions as a common primer for amplifying each base sequence to be detected. Further, the steps (2) and (3) of the first mode may be changed as follows:

(2) preparing as primers, nucleic acids each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions; and

(3) preparing a nucleic acid having a partial and sequential base sequence within the region

between a 5'-end of the A-strand and the base sequence to be detected which is located nearest the 5'-end as a primer for elongating the A-strand and preparing a nucleic acid having a base sequence complementary to a partial and sequential base sequence within the region between a 3'-end of the A-strand and the base sequence to be detected which is located nearest the 3'-end as a primer for elongating the B-strand.

That is, in addition to the B-strand elongating primer, a nucleic acid having the same base sequence as a sequential and partial base sequence which is located in the position nearer 5'-end than that of the 5'-end of the base sequence to be detected of A-strand (which is an A-strand elongating primer and a primer derived from a segment nearer the 5'-end of A-strand: herein, the phrase "derived from a position nearer the 5'-end" means "derived from a segment located in the position nearer 5'-end than that of the base sequence to be detected which is nearest the 5'-end") may be dissolved in a reaction solution to be used concomitantly as the second common primer. In some situations, the amplifying efficiency in such concomitant use may be higher.

Note that, in some situations, this method may include a step of removing a substance other than the aforementioned hybridized product (such as a template

nucleic acid, or a nucleotide monomer or enzyme used for PCR reactions) through a washing operation for removing a reaction solution on the substrate.

Those methods relate to the universal PCR as described above, and the method of detecting a nucleic acid according to a second mode of the present invention includes the above-described steps (1) to (6). This method enables the multiplex PCR.

In this case, the same detection method as that described above may be used for detecting a hybridized product. The aforementioned detection method may be used for plural single-stranded nucleic acids each having one or more partial and sequential base sequences to be detected (A-strand group: A1-strand to An-strand: $n \geq 2$). Moreover, the steps (2) and (3) according to the second mode may be changed as follows:

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions; and

(3) preparing nucleic acids each having a partial and sequential base sequence within the region between a 5'-end of each strand of the A-

strand group and the base sequence to be detected
which is located nearest the 5'-end as primers for
elongating the A-strand (PA-strand group: PA1-strand
to PAn-strand: $n \geq 2$) and preparing nucleic acids
5 each having a base sequence complementary to a
partial and sequential base sequence within the
region between a 3'-end of each strand of the A-
strand group and the base sequence to be detected
which is located nearest the 3'-end as primers for
10 elongating the B-strand (PB-strand group: PB1-strand
to PBn-strand: $n \geq 2$).

That is, in amplification reactions, plural
nucleic acids each having the same base sequence as a
sequential and partial base sequence which is located
15 in the position nearer 5'-end than that of the 5'-end
of the base sequence to be detected of the
aforementioned A-strand group (PA-strand group: PA1-
strand to PAn-strand: $n \geq 2$) may be dissolved in a
reaction solution together with the PB-strand group
20 to be used as common primers. In some situations,
the amplification efficiency in this method is higher.

The aforementioned methods enable the solid-
phase universal PCR, solid-phase multiplex PCR, or
solid-phase universal multiplex PCR using the nucleic
25 acid array.

Fig. 1 is a schematic view illustrating the
aforementioned method in the case in which A-strand

has three base sequences to be detected. In Fig. 1, a primer 1 to a primer 3 are located from the 3'-end to the 5'-end of A-strand, and on B-strand which is complementary to A-strand, a segment is defined as a common primer (univ. anti-sense primer), which is located on the segment having a sequence complementary to that of the segment selected from the region located in the position nearer 3'-end than the 3'-end of the primer 1 which is nearest the 3'-end of A-strand. Moreover, a segment is defined as a common primer (univ. sense primer), which is located on the segment having the same sequence as that in the segment selected from the region located in the position nearer 5'-end than the 5'-end of a primer 3 which is nearest the 5'-end of A-strand. The substrate has regions in which the respective primers having the same sequence as that of segments of the primer 1 to the primer 3 are immobilized, regions in which primers are immobilized are shown in Fig. 1. When PCR reactions are performed on the substrate, amplification and elongation proceed from the primers immobilized on the substrate toward the location predetermined by a common primer. In Fig. 1, at the primer-immobilized region, the 3'-end of the primer fixed at 5'-end on the substrate is elongated and amplified to the position predetermined by a common primer (univ. anti-sense primer) to form a fixed

strand corresponding to A-strand, and the fixed strand and a strand corresponding to B-strand which is amplified from B-strand may form a double-strand. By detecting the double-strand, a base sequence to be
5 detected in A-strand can be identified. When a common primer (univ. sense primer) is further added, the amplifying effect of A-strand itself may be expected, in some situations, the amplifying efficiency may be improved.

10 Next, as in the third and forth modes, in order to simplify detection steps, a labeled monomer may be utilized as a substitute for a nucleotide monomer used in PCR reactions. The situation is shown in a schematic view of Fig. 2. That is, when a labeling
15 substance is bound to a part or all of at least one of the nucleotide monomers for elongation which include bases of adenine, guanine, cytosine, and thiamine upon amplification, the aforementioned labeling substance is introduced to a nucleic acid
20 elongated from a primer immobilized on a substrate after amplification. As a result, even if steps subsequent to hybridization in the aforementioned method are omitted, detection can be performed. A labeling substance used in this method is not
25 particularly limited, but includes, for example, the above-described fluorescent dyes and radioisotopes. Also, in the case where a fluorescent dye is used for

labeling, in the detection steps in the third and forth modes, an observation step using a confocal fluorescent microscope may be additionally included.

Note that, in some situations, this method may
5 include a step of removing a substance other than a nucleic acid which is elongated and amplified by the PCR reactions and corresponds to A-strand binding to a substrate, for example, a nucleic acid which is elongated and amplified and corresponds to B-strand,
10 a template nucleic acid, a nucleotide monomer used in the PCR reactions, or an enzyme.

This method enables the solid-phase PCR capable of performing detection immediately after PCR, the solid-phase universal PCR capable of performing
15 detection immediately after PCR, the solid-phase multiplex PCR, or the solid-phase universal multiplex PCR without performing two-tiered steps in which hybridization is performed after PCR, and was required to be improved in the aforementioned
20 conventional technique.

(Other Embodiments)

An apparatus for realizing the detection methods described above fall within the scope of the present invention.

25 The above-described methods are preferred at least in points that PCR reactions and detection are performed in the form in which primer arrays are

present in the same container, which is preferred in that an apparatus or the like can be simplified and in that, as described below, PCR reactions and detection can be performed using the same method simultaneously and continuously. In such case, when the PCR reactions and detection are performed using the same method simultaneously and continuously, the process of the PCR amplification can be monitored. Accordingly, this is a preferred mode in that so-called real-time PCR is realized, that is, highly quantitative detection can be performed. An apparatus that is equipped with a PCR reaction container and detection method enabling such a detection method is a preferred mode for carrying out the present invention.

Fig. 3 illustrates a preferred embodiment of such an apparatus. In the apparatus, the PCR container comprises a substrate 1 having a surface 2 with immobilized polymers, a reaction chamber 3 and a temperature controlling unit 4. The substrate is transparent against wavelength used for detection. The reaction chamber is facing to the surface. The temperature controlling unit is placed at a position not preventing operation of detection means 5 which is placed on the side opposite to said surface in relation to said substrate.

The solution mentioned above is introduced from

an inlet port (not shown) into the reaction chamber 3 to conduct PCR reactions while the temperature is controlled by the temperature controlling unit 4 in accordance with the aforementioned schedule.

5 Fig. 4 illustrates another embodiment of such an apparatus. In the apparatus of Fig. 4, the temperature controlling unit 4 is arranged such that it surrounds also the lateral surface of the reaction container to increase the contact area.

10 Fig. 5 illustrates still another embodiment of such an apparatus. In the apparatus of Fig. 5, the entire surface except the detection area is covered by the temperature controlling unit 4. Moreover, the present invention provides a kit for detecting a
15 nucleic acid, which includes the aforementioned primer arrays, PCR reaction reagent, and detection reagent. The above-described PCR reaction container may be provided for the kit, and the PCR reaction
20 container may be formed into a cartridge form. Moreover, in the case where a nucleic acid to be detected forms a double-stranded nucleic acid, the detection reagent of the above-described kit for
25 detecting a nucleic acid is preferably a fluorescent dye as an intercalator or a groove binder, which acts on a double-stranded nucleic acid as a fluorescent dye.

EXAMPLES

Hereinafter, the present invention will be described in detail with reference to examples.

Example 1 (Solid-phase universal PCR I)

(1) Preparation of primer

5 The following seven segments (Ec1 to Ec7) of sense base sequences of genome DNA of 16s ribosomal RNA (rRNA) of *Escherischia coli* (ATCC#11775) were defined as detection targets of a solid-phase universal PCR, which were selected as primers to be
10 immobilized on solid-phase.

Ec1 5' CTCTTGCCATCGGATGTGCCCA 3' (SEQ ID NO: 1)
Ec2 5' ATACCTTGCTCATTGACGTTACCCG 3' (SEQ ID NO: 2)
Ec3 5' TTTGCTCATTGACGTTACCCGCAG 3' (SEQ ID NO: 3)
Ec4 5' ACTGGCAAGCTTGAGTCTCGTAGA 3' (SEQ ID NO: 4)
15 Ec5 5' ATACAAAGAGAAGCGACCTCGCG 3' (SEQ ID NO: 5)
Ec6 5' CGGACCTCATAAAGTGCGTCGTAGT 3' (SEQ ID NO: 6)
Ec7 5' GCGGGGAGGAAGGGAGTAAAGTTAAT 3' (SEQ ID NO: 7).

Those primers were synthesized by a synthesis company (BEX Co., Ltd.) on commission. The 5'-end of
20 each of Ec1 to Ec7 was bound to a thiol (SH) group via a linker for binding to a solid-phase substrate as described below. The Ec1 structure to which a thiol group is bound is shown below as an example.

HS(CH₂)₆OP(O₂)O-dCTCTTGCCATCGGATGTGCCCA

25 Moreover, the following base sequence (EcAP) was selected as a base sequence of a common primer of the anti-sense sequence:

EcAP 5' ATCCAACCGCAGGTTCCCCTAC 3' (SEQ ID NO: 8)

EcAP was synthesized in a general manner. All the primers were deprotected and purified based on the conventional methods.

5 (2) Extraction of Escherischia coli genome DNA

Firstly, an E.coli standard strain was cultured in accordance with the conventional method. 1.0 ml of the microorganism-cultured medium ($OD_{600} = 0.7$) was collected in a 1.5-ml microtube, and the bacterial cells were collected by centrifugation (8,500 rpm, 5 minutes, 4°C). After the supernatant was discarded, 300 μ l of Enzyme Buffer (50 mM Tris-HCl: p.H. 8.0, 25 mM EDTA) was added thereto, and the cells were resuspended using a mixer. The resuspended bacterial suspension was centrifuged again, to thereby collect bacterial cells (8,500 rpm, 5 minutes, 4°C). After the supernatant was discarded, the following enzyme solutions were added to the collected bacterial cells, and the cells were resuspended using a mixer:

20 Lysozyme 50 μ l (20 mg/ml in Enzyme Buffer)

N-Acetylmuramidase SG 50 μ l (0.2 mg/ml in Enzyme Buffer).

Next, the resuspended bacterial suspension to which the enzyme solutions were added was left to stand in an incubator at 37°C for 30 minutes, to thereby perform treatment for cell wall lysis. Subsequently, genome DNA was extracted using a kit

for purifying a nucleic acid (Mag Extractor. Genome-: manufactured by TOYOBO Co., Ltd.).

Specifically, firstly, 750 μ l of a dissolution/adsorption solution and 40 μ l of magnetic
5 beads were added to the pretreated microorganism suspension, and the mixture was stirred vigorously for 10 minutes using a tube mixer (Step 1).

The microtube was set on a separatory stand (Magical Trapper) and allowed to stand for 30 seconds,
10 to thereby collect the magnetic particles on the wall surface of the tube, and the supernatant was discarded while the tube was set on the stand (Step 2).

Next, 900 μ l of a washing solution was added
15 thereto, and the particles were resuspended by stirring using the mixer for about 5 seconds (Step 3).

Next, the microtube was set on the separatory stand (Magical Trapper) and allowed to stand for 30 seconds, to thereby collect the magnetic particles on
20 the wall surface of the tube, and the supernatant was discarded while the tube was set on the stand (Step 4).

The procedures of Steps 3 and 4 were repeated, and the second washing was performed (Step 5).
25 Subsequently, 900 μ l of 70% ethanol was added thereto, and the particles were resuspended by stirring using the mixer for about 5 seconds (Step 6).

Next, the microtube was set on the separatory stand (Magical Trapper) and allowed to stand for 30 seconds, to thereby collect the magnetic particles on the wall surface of the tube, and the supernatant was
5 discarded while the tube was set on the stand (Step 7).

The procedures of Steps 6 and 7 were repeated, and the second washing was performed with 70% ethanol (Step 8). Subsequently, 100 μ l of pure water was
10 added to the collected magnetic particles, and the mixture was stirred for 10 minutes using the tube mixer.

Next, the microtube was set on the separatory stand (Magical Trapper) and allowed to stand for 30
15 seconds, to thereby collect the magnetic particles on the wall surface of the tube, and the supernatant was collected in a new tube while the tube was set on the stand.

The collected genome DNA of *Escherischia coli*
20 was subjected to agarose electrophoresis and absorbance measurement with the wavelength of 260/280 nm in accordance with the conventional methods to assay its quality (amount of contaminated low-molecular-weight nucleic acids, the degree of
25 degradation) and collected amount. In this example, about 9 μ g of the genome DNA was collected, and degradation of the genome DNA and contamination of

rRNA were not observed. The collected genome DNA was dissolved in TE buffer so as to have a final concentration of 50 ng/ μ l, and the mixture was used in the following examples.

5 (3) Manufacture of DNA array

(3-1) Washing of glass substrate

A glass substrate made of synthetic quartz (size: 25 mm \times 75 mm \times 1 mm, manufactured by Iiyama Tokusyu Glass Co. Ltd.) was placed in a heat-resistant and alkali-resistant rack and immersed in a washing solution for ultrasonic cleaning which had been adjusted to a predetermined concentration. The substrate was immersed in the washing solution overnight, and ultrasonic cleaning was then performed for 20 minutes. Subsequently, the substrate was taken out and lightly rinsed with pure water, followed by ultrasonic cleaning in ultrapure water for 20 minutes. Next, the substrate was immersed in a 1N sodium hydroxide solution heated to 80°C for 10 minutes. Washing with pure water and washing with ultrapure water were performed again, to thereby prepare a quartz glass substrate for a DNA array.

(3-2) Surface treatment

A silane coupling agent KBM-603 (manufactured by Shin-Etsu Silicones) was dissolved in pure water so as to have a concentration of 1%, and the mixture was stirred for two hours at room temperature.

Subsequently, the glass substrate that had been previously washed was immersed in the aqueous solution of the silane coupling agent and allowed to stand for 20 minutes at room temperature. The glass substrate was drawn up, and its surface was lightly washed with pure water. Subsequently, nitrogen gas was blown on both the sides of the substrate, and the substrate was dried. Next, the treatment with the coupling agent was completed by baking the dried substrate in an oven heated to 120°C for 1 hour, to thereby introduce amino groups to the substrate surface. Next, N-(6-maleimidocaproyloxy)succinimido (hereinafter, abbreviated as EMCS; produced by Dojindo Laboratories) was dissolved in a mixed solvent of dimethylsulfoxide and ethanol (1:1) so as to have a final concentration of 0.3 mg/ml to prepare an EMCS solution. The baked glass substrate was left to cool and immersed in the prepared EMCS solution at room temperature for 2 hours. Through such treatment, the amino groups introduced onto the surface by using the silane coupling agent reacted with succinimido groups of EMCS, to thereby introduce maleimide groups onto the surface of the glass substrate. The glass substrate was drawn up from the EMCS solution, and the substrate was washed with the above-described mixed solvent and further washed with ethanol, followed by drying under nitrogen gas atmosphere.

(3-3) Primers

The primers to be immobilized on solid-phase (Ec1 to Ec7) described in (1) were separately dissolved in pure water, and each mixture was
5 dispensed so as to have a final concentration (when it was dissolved as described below) of 10 μ M, followed by freeze-drying, to thereby remove water.

(3-4) Discharge of primer DNA by BJ printer, and binding to substrate

10 An aqueous solution (mixed solvent) containing 7.5 wt% of glycerin, 7.5 wt% of thiodiglycol, 7.5 wt% of urea, and 1.0 wt% of Acetylenol EH (manufactured by Kawaken Fine Chemicals Co., Ltd.) was prepared. Subsequently, each of previously prepared 7 primers
15 was dissolved in the aforementioned mixed solvent so as to have a prescribed concentration. The resultant DNA solution was filled into an ink tank for a bubble jet printer (trade name: BJF-850, manufactured by Canon Inc.), and the tank was attached to a print
20 head.

Note that the bubble jet printer used herein had been modified so as to enable print on a flat plate. Moreover, the bubble jet printer can perform spotting of about 5 pl of a DNA solution at a pitch
25 of about 120 μ m by inputting a printing pattern in accordance with a predetermined file creation method.

Subsequently, a printing operation for one

glass substrate was performed using the modified bubble jet printer for preparing a DNA array. After confirming that printing was performed without fail, the array was allowed to stand in a humidified chamber for 30 minutes to react maleimide groups on the surface of the glass substrate with thiol groups at the end of the nucleic acid primer.

(3-5) Washing

After the reaction was performed for 30 minutes, the DNA solution remaining on the surface was washed out with 10 mM of phosphate buffer (pH 7.0) containing 100 mM of NaCl, to thereby yield a DNA array in which single-stranded DNA was immobilized on the surface of the glass substrate.

(4) Solid-phase PCR

A method of amplifying *Escherichia coli* genome DNA was described below.

Premix PCR reagent (TAKARA ExTaq)	25 μ l
Template <i>Escherichia coli</i> genome DNA	2 μ l (100 ng)
Primer EcAP	2 μ l (20 pmole)
H ₂ O	21 μ l
<hr/>	
Total	50 μ l

A microchamber in which a primer-binding site of the above-described primer array was covered and the temperature can be controlled was manufactured,

and a reaction solution of the aforementioned composition was sealed in the chamber. Then, amplification reactions were performed in accordance with the following protocol. More specifically,

5 after an incubation step at 95°C/10 minutes, a denaturation step at 92°C/45 seconds, an annealing step at 55°C/45 seconds, and an elongation step at 72°C/45 seconds were defined as one cycle, and the cycle was repeated 35 times, and finally an

10 incubation step at 72°C/10 minutes was performed.

	95°C	10 minutes	
	92°C	45 seconds	←
	55°C	45 seconds	35 Cycles
15	72°C	45 seconds	—
	72°C	10 minutes	

After the completion of the reactions, 1 µl of a solution prepared by previously diluting a dye of which (fluorescence is enhanced under coexistence of

20 a double-stranded nucleic acid, SYBR (registered trademark) Green I (Molecular Probes: trade name SYBR (registered trademark) Green I nucleic acid gel stain 10,000 × concentrate in DMSO) to 200-fold with pure

25 water was added to the chamber, and the primer array was placed under the conditions of 65°C/3 minutes → 92°C/2 minutes → 45°C/3 hours for hybridization.

Next, the primer array was washed under the conditions of $2 \times \text{SSC}/0.1\% \text{ SDS}$ (25°C)/2 minutes $\rightarrow 2 \times \text{SSC}$ (20°C)/2 minutes \rightarrow pure water (10°C)/2 minutes, and the array was taken out of the chamber and dried.

5 (5) Fluorescence measurement

The DNA array after the completion of the hybridization reaction was subjected to fluorescence measurement using a fluorescence detecting apparatus for DNA array (manufactured by Axon Instruments, GenePix 4000B) (excitation wavelength: 532 nm, photomultiplier voltage: 400 V). The measurement results are shown in Table 1. Note that, in this example, all the operations are performed twice, so that the results were separately shown in Table 1.

15

Table 1

Primer No.	Fluorescence luminance	
	First time	Second time
Ec1	8,700	8,500
Ec2	9,100	9,050
Ec3	7,750	7,650
Ec4	11,000	11,000
Ec5	10,400	10,550
Ec6	7,400	7,400
Ec7	5,900	6,000

The numeric values of the fluorescent luminance in Table 1 represent pixel average luminance

(resolution: 5 μ m). As is clear from Table 1, the results obtained by amplifying genome DNA extracted from *Escherischia coli* by the solid-phase universal PCR can be detected with high reproducibility.

5 Example 2 (Solid-phase universal PCR II)

The solid-phase PCR and fluorescence detection were performed in accordance with the same method as that in Example 1 except that, in addition to the primers used for the solid-phase PCR in Example 1, a
10 common primer (EcSP) of the sense sequence having the following base sequence was used at the same concentration as that of a common primer of the anti-sense sequence.

EcSP 5' GCGGCAGGCCTAACACATGCAAG 3' (SEQ ID NO: 9).

15 In Example 2, when the solid-phase PCR cycle was repeated 31 times, substantially the same results as those in Example 1 were obtained. The results reveal that the efficiency of the solid-phase PCR could be improved by letting both the common primers
20 of the sense sequence and anti-sense sequence coexist in a solution.

Example 3 (Solid-phase multiplex universal PCR I)

(1) Preparation of primers

In this example, genome DNA of rRNA of
25 *Pseudomonas aeruginosa* (ATCC#10145) will be detected simultaneously with *Escherischia coli* detected in Example 1. For this purpose, firstly, 8 thiolized

primers of the sense sequence for *Pseudomonas aeruginosa* (Pa1 to Pa8) were synthesized in the same way as that in Example 1. The base sequences of the primers are shown below.

5 Pa1 5' TGAGGGAGAAAGTGGGGGATCTTC 3' (SEQ ID NO: 10)
Pa2 5' TCAGATGAGCCTAGGTTCGATTAGC 3' (SEQ ID NO: 11)
Pa3 5' GAGCTAGAGTACGGTAGAGGGTGG 3' (SEQ ID NO: 12)
Pa4 5' GTACGGTAGAGGGTGGTGAATTTTC 3' (SEQ ID NO: 13)
Pa5 5' GACCACCTGGACTGATACTGACAC 3' (SEQ ID NO: 14)
10 Pa6 5' TGGCCTTGACATGCTGAGAACTTTC 3' (SEQ ID NO: 15)
Pa7 5' TTAGTTACCAGCACCTCGGGTGG 3' (SEQ ID NO: 16)
Pa8 5' TAGTCTAACCGCAAGGGGGACG 3' (SEQ ID NO: 17).

Moreover, the following base sequence (PaAP) was selected as a base sequence of a common primer of the anti-sense sequence:

15 PaAP 5' ATCCAGCCGCAGGTTCCCCTAC 3' (SEQ ID NO: 18).

(2) Fluorescence detection

Extraction of *Pseudomonas aeruginosa* genome, manufacture of a DNA microarray immobilized with
20 *Escherichia coli* primers and *Pseudomonas aeruginosa* primers, solid-phase PCR (*Escherichia coli* genome DNA and *Pseudomonas aeruginosa* genome DNA: 10 ng each, EcAP and PaAP: 20 pmole each), hybridization, washing, etc. were performed in the same way as that in
25 Example 1, and fluorescence detection was performed. The results are shown in Table 2.

Table 2

Primer No.	Fluorescence luminance	
	Escherischia coli	Pseudomonas aeruginosa
Pa1	5,800	300
Pa2	6,050	550
Pa3	5,150	1,700
Pa4	7,400	650
Pa5	6,900	1,600
Pa6	4,900	2,650
Pa7	3,900	600
Pa8	-	350

Table 2 shows that all the given primer sequences of rRNA genomes of both *Escherischia coli* and *Pseudomonas aeruginosa* were simultaneously detected by the solid-phase multiplex universal PCR. Example 4 (Solid-phase multiplex universal PCR II)

The solid-phase PCR and fluorescence detection were performed in the same manner as that in Example 3 except that, in addition to the primers used for the solid-phase PCR in Example 2, common primers of the respective sense sequences of *Escherischia coli* and *Pseudomonas aeruginosa* (EcSP and PaSP) were used at the same concentration. The base sequence of PaSP is shown below.

PaSP 5' GCGGCAGGCTTAACACATGCAAG 3' (SEQ ID NO: 19).

In Example 4, under the condition that the

cycle of the solid-phase PCR was repeated at given times fewer than those in Example 3 by about one or more, almost the same results as those in Example 3 were obtained. The results reveal that, in the solid-phase multiplex universal PCR, the efficiency of the solid-phase PCR could be improved by letting both the common primers of the sense sequence and anti-sense sequence coexist in a solution.

Example 5 (Incorporated-label solid-phase universal PCR)

Extraction of *Escherichia coli* genome, synthesis of primers, and manufacture of a primer array were performed in completely the same manner as that in Example 1, and the solid-phase PCR was then performed using a PCR solution of the following composition.

Premix PCR reagent (TAKARA ExTaq)	25 μ l
Template <i>Escherichia coli</i> genome DNA	2 μ l (100 ng)
Primer EcAP	2 μ l (20 pmole)
Cy-3 dUTP (Amersham Biosciences K.K.: 1 mM)	2 μ l (2 nmol)
H ₂ O	19 μ l
Total	50 μ l

Subsequently, the primer array was washed, without keeping the same under to hybridization

conditions, under the conditions of 2 × SSC/0.1% SDS
(92°C)/2 minutes → 2 × SSC/0.1% SDS (92°C)/2 minutes
→ 2 × SSC/0.1% SDS (25°C)/2 minutes → 2 × SSC
(20°C)/2 minutes → pure water (20°C)/2 minutes, and
5 the array was taken out of the chamber and dried.

Subsequently, fluorescence detection was
performed in the same manner as that in Example 1.
The results are shown in Table 3.

10

Table 3

Primer No.	Escherischia coli
	Fluorescence luminance
Ec1	11,500
Ec2	11,900
Ec3	10,000
Ec4	14,800
Ec5	13,200
Ec6	9,500
Ec7	7,700

In accordance with the detection method of the
present invention, fluorescence detection was
performed adopting an incorporated label in the
15 solid-phase universal PCR. As a result, fluorescence
detection was easily accomplished as shown in Table 3.
Accordingly, the results reveal that fluorescence
detection can be performed without requiring the
hybridization step.

Example 6

The solid-phase universal PCR or solid-phase multiplex universal PCR in Examples 2 to 4 was performed using the incorporated label in Example 5.

5 Although the respective fluorescence intensities were different from those in Examples 2 to 4, the results reveal that the rRNA genome DNA of *Escherischia coli* or the rRNA genome DNAs of both *Escherischia coli* and *Pseudomonas aeruginosa* can be detected simultaneously.

10 Example 7 (Common primer-labeled solid-phase universal PCR)

Detection of *Escherischia coli* genome DNA was performed by the solid-phase universal method in completely the same manner as that in Example 1
15 except that the common primer EcAP was labeled with tetramethylrhodamine, hybridization was performed without coexistence with a fluorescent dye such as SYBR green, and fluorescence detection was performed using the aforementioned tetramethylrhodamine. The
20 structure of tetramethylrhodamine-labeled EcAP is shown below.

5' Rho-CONH(CH₂)₆OP(O₂)O-d ATCCAACCGCAGGTTCCCCTAC 3'
(Rho = tetramethylrhodamine).

As a result, almost the same results as those
25 in Example 1 were obtained although the differences in fluorescence intensities were observed.

A method of labeling the common primer of anti-

sense sequence was performed in the methods of Examples 2 to 4. As a result, the results confirm that all the methods were effective were obtained.

5 The present invention is not limited to the above embodiments and various changes and modifications can be made within the spirit and scope of the present invention. Therefore to apprise the public of the scope of the present invention, the following claims are made.

10

This application claims priority from Japanese Patent Application No. 2004-070986 filed on March 12, 2004, which is hereby incorporated by reference
15 herein.

CLAIMS

1. A method of detecting a nucleic acid,
comprising the steps of:

5 (1) preparing a single-stranded nucleic acid
having plural partial and sequential base sequences
to be detected (A-strand) and a single-stranded
nucleic acid having a base sequence complementary to
a base sequence of the A-strand (B-strand);

10 (2) preparing nucleic acids as primers each
having one of the plural base sequences to be
detected, immobilizing the respective primers
independently in separate regions on a substrate, and
preparing a primer array in which the respective base
sequences to be detected are distributed in the
15 primer-immobilized regions;

(3) preparing a nucleic acid having a sequence
complementary to a partial and sequential base
sequence within the region between a 3'-end of the A-
strand and the base sequence to be detected which is
20 located nearest the 3'-end as a primer for elongating
the B-strand;

(4) performing PCR reactions using the A-strand
and B-strand as templates, and using the primers
immobilized on the substrate, and the primer for
25 elongating the B-strand;

(5) forming a hybridized product of a nucleic
acid corresponding to the A-strand which has been

elongated and amplified as a result of the PCR reactions and bound to the substrate and a nucleic acid corresponding to the B-strand which has been elongated and amplified and has not bound to the substrate; and

(6) detecting the base sequence to be detected by detecting the hybridized product in the respective primer-immobilized regions in the array.

2. A method of detecting a nucleic acid, comprising the steps of:

(1) preparing a single-stranded nucleic acid having plural partial and sequential base sequences to be detected (A-strand) and a single-stranded nucleic acid having a base sequence complementary to a base sequence of the A-strand (B-strand);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions;

(3) preparing a nucleic acid having a partial and sequential base sequence within the region between a 5'-end of the A-strand and the base sequence to be detected which is located nearest the 5'-end as a primer for elongating the A-strand and

preparing a nucleic acid having a sequence
complementary to a partial and sequential base
sequence within the region between a 3'-end of the A-
strand and the base sequence to be detected which is
5 located nearest the 3'-end as a primer for elongating
the B-strand;

(4) performing PCR reactions using the A-strand
and B-strand as templates, and using the primers
immobilized on the substrate, the primer for
10 elongating the A-strand, and the primer for
elongating the B-strand;

(5) forming a hybridized product of a nucleic
acid corresponding to the A-strand which has been
elongated and amplified as a result of the PCR
15 reactions and bound to the substrate and a nucleic
acid corresponding to the B-strand which has been
elongated and amplified and has not bound to the
substrate; and

(6) detecting the base sequence to be detected
20 by detecting the hybridized product in the respective
primer-immobilized regions in the array.

3. A method of detecting a nucleic acid,
comprising the steps of:

(1) preparing plural single-stranded nucleic
25 acids each having a partial and sequential base
sequence to be detected (A-strand group: A1-strand to
An-strand: $n \geq 2$) and a group of single-stranded

nucleic acids each having a base sequence complementary to a base sequence of each strand of the A-strand group (B-strand group: B1-strand to Bn-strand: $n \geq 2$);

5 (2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base
10 sequences to be detected are distributed in the primer-immobilized regions;

 (3) preparing nucleic acids each having a sequence complementary to a partial and sequential base sequence within the region between a 3'-end of
15 each strand of the A-strand group and the base sequence to be detected which is located nearest the 3'-end as primers for elongating the B-strands (PB-strand group: PB1-strand to PBn-strand: $n \geq 2$);

 (4) performing PCR reactions using each strand
20 of the A-strand group and each corresponding strand of B-strand group as templates, and using the primers immobilized on the substrate, and the plural primers for elongating the B-strands of the PB-strand group;

 (5) forming a hybridized product of a nucleic
25 acid corresponding to the A-strand group which has been elongated and amplified as a result of the PCR reactions and bound to the substrate and a nucleic

acid corresponding to the B-strand group which has been elongated and amplified and has not bound to the substrate; and

(6) detecting the base sequence to be detected
5 by detecting the hybridized product in the respective primer-immobilized regions in the array.

4. A method of detecting a nucleic acid,
comprising the steps of:

(1) preparing plural single-stranded nucleic
10 acids each having a partial and sequential base sequence to be detected (A-strand group: A1-strand to An-strand: $n \geq 2$) and a group of single-stranded nucleic acids each having a base sequence complementary to a base sequence of each strand of
15 the A-strand group (B-strand group: B1-strand to Bn-strand: $n \geq 2$);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers
20 independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions;

(3) preparing nucleic acids each having a
25 partial and sequential base sequence within the region between a 5'-end of each strand of the A-strand group and the base sequence to be detected

which is located nearest the 5'-end as primers for
elongating the A-strands (PA-strand group: PA1-strand
to PAn-strand: $n \geq 2$) and preparing nucleic acids
having a sequence complementary to a partial and
5 sequential base sequence within the region between a
3'-end of each strand of the A-strand group and the
base sequence to be detected which is located nearest
the 3'-end as primers for elongating the B-strands
(PB-strand group: PB1-strand to PBn-strand: $n \geq 2$);
10 (4) performing PCR reactions using each strand
of the A-strand group and each corresponding strand
of the B-strand group as templates, and using the
primers immobilized on the substrate, the primers for
elongating the A-strands of the PA-strand group, and
15 the primer for elongating the B-strand of the PB-
strand group;
(5) forming a hybridized product of a nucleic
acid corresponding to the A-strand group which has
been elongated and amplified as a result of the PCR
20 reactions and bound to the substrate and a nucleic
acid corresponding to the B-strand group which has
been elongated and amplified and has not bound to the
substrate; and
(6) detecting the base sequence to be detected
25 by detecting the hybridized product in the respective
primer-immobilized regions in the array.

5. A method of detecting a nucleic acid

according to any one of claims 1 to 4, further comprising a step of washing and removing a reaction solution on the substrate after the PCR reactions.

6. A method of detecting a nucleic acid
5 according to any one of claims 1 to 4, wherein the primer for elongating the B-strand is labeled, and the hybridized product is detected using the label.

7. A method of detecting a nucleic acid
according to claim 5, wherein the label is a
10 fluorescent dye.

8. A method of detecting a nucleic acid
according to claim 7, further comprising a step of observing the fluorescent dye using a confocal fluorescent microscope for detecting the hybridized
15 product.

9. A method of detecting a nucleic acid
according to any one of claims 1 to 4, wherein the hybridized product is detected using a fluorescent dye as an intercalator or a groove binder which
20 interacts with a double-stranded nucleic acid.

10. A method of detecting a nucleic acid
according to claim 9, further comprising a step of observing the fluorescent dye using a confocal fluorescent microscope for detecting the hybridized
25 product.

11. A method of detecting a nucleic acid,
comprising the steps of:

(1) preparing a single-stranded nucleic acid having plural partial and sequential base sequences to be detected (A-strand) and a single-stranded nucleic acid having a base sequence complementary to a base sequence of the A-strand (B-strand);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions;

(3) preparing a nucleic acid having a sequence complementary to a partial and sequential base sequence within the region between a 3'-end of the A-strand and the base sequence to be detected which is located nearest the 3'-end as a primer for elongating the B-strand;

(4) performing PCR reactions using the A-strand and the B-strand as templates, and using the primers immobilized on the substrate, and the primer for elongating the B-strand, and nucleotide monomers with a part or all of at least one group of the nucleotide monomer being labeled; and

(5) detecting a nucleic acid corresponding to the A-strand which has been elongated and amplified from a primer binding to the substrate via the label

incorporated in the nucleic acid.

12. A method of detecting a nucleic acid,
comprising the steps of:

(1) preparing a single-stranded nucleic acid
5 having plural partial and sequential base sequences
to be detected (A-strand) and a single-stranded
nucleic acid having a base sequence complementary to
a base sequence of the A-strand (B-strand);

(2) preparing nucleic acids as primers each
10 having one of the plural base sequences to be
detected, immobilizing the respective primers
independently in separate regions on a substrate, and
preparing a primer array in which the respective base
sequences to be detected are distributed in the
15 primer-immobilized regions;

(3) preparing a nucleic acid having a partial
and sequential base sequence within the region
between a 5'-end of the A-strand and the base
sequence to be detected which is located nearest the
20 5'-end as a primer for elongating the A-strand and
preparing a nucleic acid having a base sequence
complementary to a partial and sequential base
sequence within the region between a 3'-end of the B-
strand and the base sequence to be detected which is
25 located nearest the 3'-end as a primer for elongating
the B-strand;

(4) performing PCR reactions using the A-strand

and the B-strand as templates, and using the primers immobilized on the substrate, the primer for elongating the A-strand, and the primer for elongating the B-strand, and nucleotide monomers with a part or all of at least one group of the nucleotide monomer being labeled; and

(5) detecting a nucleic acid corresponding to the A-strand which has been elongated and amplified from a primer binding to the substrate via the label incorporated in the nucleic acid.

13. A method of detecting a nucleic acid, comprising the steps of:

(1) preparing plural single-stranded nucleic acids each having a partial and sequential base sequence to be detected (A-strand group: A1-strand to An-strand: $n \geq 2$) and a group of single-stranded nucleic acids each having a base sequence complementary to a base sequence of each strand of the A-strand group (B-strand group: B1-strand to Bn-strand: $n \geq 2$);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions;

(3) preparing nucleic acids each having a sequence complementary to a partial and sequential base sequence within a region between a 3'-end of each strand of the A-strand group and the base sequence to be detected which is located nearest the 3'-end as primers for elongating the B-strands (PB-strand group: PB1-strand to PBn-strand: $n \geq 2$);

(4) performing PCR reactions using each strand of the A-strand group and each corresponding strand of the B-strand group as templates, and using the primers immobilized on the substrate and the plural primers for elongating the B-strands of the PB-strand group, and nucleotide monomers with a part or all of at least one group of the nucleotide monomer being labeled; and

(5) detecting a nucleic acid corresponding to the A-strand which has been elongated and amplified from a primer binding to the substrate via the label incorporated in the nucleic acid.

14. A method of detecting a nucleic acid, comprising the steps of:

(1) preparing plural single-stranded nucleic acids each having a partial and sequential base sequence to be detected (A-strand group: A1-strand to An-strand: $n \geq 2$) and a group of single-stranded nucleic acids each having a base sequence complementary to a base sequence of each strand of

the A-strand group (B-strand group: B1-strand to Bn-strand: $n \geq 2$);

(2) preparing nucleic acids as primers each having one of the plural base sequences to be
5 detected, immobilizing the respective primers independently in separate regions on a substrate, and preparing a primer array in which the respective base sequences to be detected are distributed in the primer-immobilized regions;

10 (3) preparing nucleic acids each having a partial and sequential base sequence within the region between a 5'-end of each strand of the A-strand group and the base sequence to be detected which is located nearest the 5'-end as primers for
15 elongating the A-strands (PA-strand group: PA1-strand to PAn-strand: $n \geq 2$) and preparing nucleic acids each having a base sequence complementary to a partial and sequential base sequence within the region between a 3'-end of each strand of the A-
20 strand group and the base sequence to be detected which is located nearest the 3'-end as primers for elongating the B-strand (PB-strand group: PB1-strand to PBn-strand: $n \geq 2$);

(4) performing PCR reactions using each strand
25 of the A-strand group and each corresponding strand of the B-strand group as templates, and using the primers immobilized on the substrate and respective

primers of the PA-strand group and PB-strand group, and nucleotide monomers with a part or all of at least one group of the nucleotide monomer being labeled; and

5 (5) detecting a nucleic acid corresponding to the A-strand which has been elongated and amplified from a primer binding to the substrate via the label incorporated in the nucleic acid.

15 15. A method of detecting a nucleic acid according to any one of claims 11 to 14, further comprising a step of washing and removing a reaction solution on the substrate after the PCR reactions.

16. A method of quantitative determination of a nucleic acid based on signals detected according to
15 any one of claims 1 to 4 and 11 to 14.

17. A method of detecting a nucleic acid according to any one of claims 11 to 14, wherein the label is a fluorescent dye.

18. A method of detecting a nucleic acid
20 according to claim 17, further comprising a step of observing the fluorescent dye using a confocal fluorescent microscope for detecting the hybridized product.

19. A method of detecting a nucleic acid
25 according to any one of claims 1 to 4 and 11 to 14, wherein at least the PCR reactions and nucleic acid detections are performed in a form in which the

primer arrays are present in the same container.

20. A method of detecting a nucleic acid according to claim 19, wherein the respective PCR reactions and nucleic acid detections are performed while observing intermittently using the same means.

21. An apparatus for detecting a nucleic acid, which enables the method of detecting a nucleic acid according to claim 19, comprising:

a PCR reaction container; and
detection means.

22. An apparatus for detecting a nucleic acid according to claim 21,

wherein said PCR container comprises a substrate having a surface with immobilized polymers, a reaction chamber and a temperature controlling unit,

wherein said substrate is transparent against wavelength used for detection

wherein said reaction chamber is facing to said surface,

wherein said temperature controlling unit is placed at a position not preventing operation of said detection means, and

wherein said detection means is placed on the side opposite to said surface in relation to said substrate.

23. A kit for detecting a nucleic acid, comprising a primer array; a PCR reaction reagent;



and a nucleic acid detecting reagent, for performing the method according to any one of claims 1 to 4 and 11 to 14.

24. A kit for detecting a nucleic acid
5 according to claim 23, wherein the nucleic acid detecting reagent is a fluorescent dye serving as an intercalator or groove binder which acts on a double-stranded nucleic acid.

1 / 3


FIG. 1

GENE (TEMPLATES) AND PRIMER ARRAY






2/3

FIG. 2



3 / 3

FIG. 3**FIG. 4****FIG. 5**

SEQUENCE LISTING

<110> Canon Kabushiki Kaisha

<120> Method of detecting nucleic acid

<130> 10009684W001

<150> JP2004-070986

<151> 2004-03-12

<160> 19

<170> PatentIn version 3.3

<210> 1

<211> 22

<212> DNA

<213> Synthesized

<400> 1

ctcttgccat cggatgtgcc ca

22

<210> 2

<211> 26

<212> DNA

<213> Synthesized

<400> 2

atacctttgc tcattgacgt tacccg

26

<210> 3

<211> 24

<212> DNA

<213> Synthesized

<400> 3

tttgctcatt gacgttaccc gcag

24

<210> 4

<211> 24

<212> DNA

<213> Synthesized

<400> 4

actggcaagc ttgagtctcg taga

24

<210> 5

<211> 23

<212> DNA

<213> Synthesized

<400> 5

atacaaagag aagcgacctc gcg

23

<210> 6

<211> 25

<212> DNA

<213> Synthesized

<400> 6

cggacctcat aaagtgcgtc gtagt

25

2/3

<210> 7
<211> 26
<212> DNA
<213> Synthesized

<400> 7
gcggggagga agggagtaaa gttaat

26

<210> 8
<211> 22
<212> DNA
<213> Synthesized

<400> 8
atccaaccgc aggttcccct ac

22

<210> 9
<211> 23
<212> DNA
<213> Synthesized

<400> 9
gcggcaggcc taacacatgc aag

23

<210> 10
<211> 24
<212> DNA
<213> Synthesized

<400> 10
tgaggagaaa agtgggggat ctic

24

<210> 11
<211> 25
<212> DNA
<213> Synthesized

<400> 11
tcagatgagc ctaggtcgga ttagc

25

<210> 12
<211> 24
<212> DNA
<213> Synthesized

<400> 12
gagctagagt acggtagagg gtgg

24

<210> 13
<211> 25
<212> DNA
<213> Synthesized

<400> 13
gtacggtaga gggtagtgga atttc

25

<210> 14
<211> 24
<212> DNA
<213> Synthesized

<400> 14

3/3

gaccacctgg actgatactg acac

24

<210> 15
<211> 25
<212> DNA
<213> Synthesized

<400> 15
tggccttgac atgctgagaa ctttc

25

<210> 16
<211> 23
<212> DNA
<213> Synthesized

<400> 16
ttagttacca gcacctcggg tgg

23

<210> 17
<211> 22
<212> DNA
<213> Synthesized

<400> 17
tagtctaacc gcaaggggga cg

22

<210> 18
<211> 22
<212> DNA
<213> Synthesized

<400> 18
atccagccgc aggttcccct ac

22

<210> 19
<211> 23
<212> DNA
<213> Synthesized

<400> 19
gcggcaggct taacacatgc aag

23

INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP2005/004881

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01/71041 A (MERGEN LTD; YU, ZAILIN; PENG, ZAOYUAN; HU, QIANJIN) 27 September 2001 (2001-09-27) the whole document -----	1,3, 5-11,13, 16-23
X	WO 02/081743 A (HAMILL, BRENDAN) 17 October 2002 (2002-10-17) the whole document -----	1-24
X	WO 01/48242 A (MERGEN LTD; HU, QIANJIN; PENG, ZAOYUAN; YU, ZAILIN) 5 July 2001 (2001-07-05) abstract; figure 2 page 14 page 19, paragraph 90 ----- -/-	11,13, 15-24

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

20 July 2005

Date of mailing of the international search report

27/07/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Cornelis, K

INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP2005/004881

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2004/017068 A (HITACHI HIGH TECH CORP 'JP!; MATSUI TAKUYA 'JP!; MIYAHARA YUJI 'JP!; H) 26 February 2004 (2004-02-26) the whole document -----	1-24
A	PATENT ABSTRACTS OF JAPAN vol. 2002, no. 02, 2 April 2002 (2002-04-02) & JP 2001 299346 A (NABA HIROYUKI), 30 October 2001 (2001-10-30) abstract -----	1-24
A	WO 03/019189 A (CAO, WEI; LI, WENQUAN; ZHANG, YUEJIAN; SONG, JUN) 6 March 2003 (2003-03-06) the whole document -----	1-24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP2005/004881

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0171041	A	27-09-2001	US 6376191 B1 AU 5093001 A CA 2374406 A1 CN 1426481 A ,C EP 1301623 A2 JP 2003527867 T US 2002086322 A1 WO 0171041 A2	23-04-2002 03-10-2001 27-09-2001 25-06-2003 16-04-2003 24-09-2003 04-07-2002 27-09-2001
WO 02081743	A	17-10-2002	CA 2442939 A1 EP 1373566 A2 WO 02081743 A2 GB 2377220 A ,B JP 2004525640 T	17-10-2002 02-01-2004 17-10-2002 08-01-2003 26-08-2004
WO 0148242	A	05-07-2001	AU 2282801 A CN 1500150 A EP 1255860 A2 JP 2003523183 T WO 0148242 A2 US 2001036632 A1	09-07-2001 26-05-2004 13-11-2002 05-08-2003 05-07-2001 01-11-2001
WO 2004017068	A	26-02-2004	WO 2004017068 A1 EP 1542009 A1	26-02-2004 15-06-2005
JP 2001299346	A	30-10-2001	NONE	
WO 03019189	A	06-03-2003	CN 1407113 A WO 03019189 A1	02-04-2003 06-03-2003